Dosing for Infants and Children

TYLENOL'

TYLENOL'

From Your Healthcare Professional

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS. Infants' TYLENOL® Children's TYLENOL® If possible, use TYLENOL **Oral Suspension** weight to dose; **Oral Suspension** otherwise use age. **Active Ingredient:** Active Ingredient: Acetaminophen 160 mg Acetaminophen 160 mg mL = milliliter (in each 5 mL) Available in: (in each 5 mL or 1 tsp) Available in: tsp = teaspoon Grape Grape Use only as directed. Use only as directed. WEIGHT AGE Cherry 6-11 lbs 0-3 mos 1.25 mL 교 3.75 교 25 교 25 교 25 12-17 lbs 2.5 mL 4-11 mos 교 3.75 교 3.75 교 3.75 교 3.75 18-23 lbs 12-23 mos 3.75 mL 11.25 파리 3.75 파리 5 24-35 lbs 5 mL 2-3 yrs 5 型 5 型 3.75 5 mL (1 tsp) mL 36-47 lbs 4-5 yrs 7.5 7.5 mL (1½ tsp) mL 48-59 lbs 6-8 yrs 10 10 mL (2 tsp) mL 60-71 lbs 9-10 yrs 12.5 12.5 mL (2½ tsp) mL

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date:

11 yrs

72-95 lbs

This dosing recommendation from your doctor will expire in 14 DAYS.

© Johnson & Johnson Consumer Inc. 2015 • CTCM-0387

- Read and follow the label on all TYLENOL® products
- Repeat dose every 4 hours while symptoms last
- Do NOT give more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device (syringe or dosing cup) that came with the product

15 mL (3 tsp)

15 mL

 All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)